

AGE AND GROWTH OF PIKE PERCH SANDER *LUCIOPERCA* (PERCIDAE) IN THE GHRIB RESERVOIR (NORTHWEST ALGERIA)Amina BOUAMRA^{1*}, Billel BELAIFA¹, Lamya CHAOU², Mohamed Hichem KARA²
& Abdeslem ARAB¹¹ Dynamic and Biodiversity Laboratory, Faculty of Biological Science, U.S.T.H.B, BP 32, El Alia, Algiers, Algeria. E-mails: belaifa.billel@outlook.com, abdeslama@yahoo.fr² Marine Bioresources Laboratory, University of Annaba, Badji-Mokhtar, Algeria. E-mails: chaouilamya@hotmail.com, kara_hichem@yahoo.com* Corresponding author: Tel: + 213 661530004; Fax: +213 21247217; E-mail: bouamra.amina@outlook.com

RESUME.— *Âge et croissance du Sandre Sander lucioperca* (Percidae) dans le réservoir de Ghrib (Nord-Ouest Algérien).— L'objectif de cette étude était de déterminer les paramètres de base spécifiques aux populations nécessaires à l'évaluation des stocks de poissons dans le barrage de Ghrib et de les comparer avec des données provenant d'autres régions. L'âge et la croissance du Sandre, *Sander lucioperca* (Linnaeus, 1758), ont été étudiés mensuellement de mai 2013 à avril 2014 dans ce barrage algérien, à partir de 849 spécimens. En utilisant la méthode scalimétrique, nous avons pu identifier 9 groupes d'âge dans la population échantillonnée ($21 < L_t < 88,2$ cm). Les relations taille-poids entre la longueur totale (L_t) et le poids (W) ont été établies pour la masse brute et la masse éviscérée, sexes séparés et sexes confondus, soit dans ce dernier cas : $W_t = 0,0033 L_t^{3,237}$ ($77 \leq W_t \leq 7245$ g) et $W_e = 0,003 L_t^{3,271}$ ($59 \leq W_e \leq 6229$ g). La taille du poisson au moment de l'apparition des écailles a été estimée à $L_{40} = 78,86$ mm. Les valeurs moyennes des tailles de poisson obtenues par rétro-calcul à chaque formation des anneaux s'adaptent bien au modèle de croissance de Von Bertalanffy : $L_t = 125,72 (1 - e^{-0,13(t + 1,29)})$. L'indice de performance de croissance (ϕ) est de 3,31. Les paramètres de croissance déterminés par la méthode de Bhattacharya s'ajustent bien au modèle de croissance de Von Bertalanffy : $L_t = 132,17 (1 - e^{-0,12(t + 1,69)})$. Le test χ^2 appliqué aux résultats des deux méthodes n'a pas montré de différence significative.

SUMMARY.— Our objective was to estimate growth parameters for Pike perch *Sander lucioperca* (Linnaeus, 1758) in the Ghrib reservoir. Additionally, we compared these values with those from populations in other regions. We collected 849 pike perches ($21 < TL < 88.2$ cm) from May 2013 to April 2014. Direct scale readings were performed by counting the number of rings, and the results were compared with data obtained by back-calculating the length at different ages. The individuals in the sample were aged between 1⁺ and 9⁺. Size-mass relationships were calculated for the total mass and the eviscerated mass: $TW = 0.0033 TL^{3,237}$ ($77 \leq TW \leq 7245$ g) and $EW = 0.003 TL^{3,271}$ ($59 \leq EW \leq 6229$ g). The mean size of fish at the time of first scale formation was estimated as $L_{40} = 78.86$ mm. The median values of the fish sizes obtained by retro-calculation were consistent with the Von Bertalanffy model of growth: $TL = 125.72 (1 - e^{-0.13(t + 1.29)})$. The growth performance index (ϕ) was 3.31. The parameters of growth determined by the Bhattacharya's method were consistent with the Von Bertalanffy growth model: $TL = 132.17 (1 - e^{-0.12(t + 1.69)})$. The χ^2 test applied to the results of the two methods did not show any significant difference.

In Algeria, Percidae are represented by two species: *Sander lucioperca* (Linnaeus, 1758) and *Perca fluviatilis* (Linnaeus, 1758). *Sander lucioperca* is a European species that was historically distributed from Elba and the Baltic Sea in the east to South-West Russia in the west (Deelder & Willemsen, 1964; Sonesten, 1991; Campbell, 1992). Currently, the species is found in the area from the Iberian peninsula to the Sea of Aral and from Scandinavia to the Maghreb countries. Additionally, it has been introduced in the Azores and into certain lakes of the United States (Poulet, 2004).

In 1985, specimens from Hungary were introduced to Lake Oubéira and reservoirs in the north of Algeria, including Ghrib reservoir, to supplement the native fish fauna and aid development of aquaculture (Meddour *et al.*, 2005; Kara, 2012). The species currently supports a robust and economically important fishery for individual and commercial fishers in the region.

However, introduction of pike perch has resulted in conflict in some areas of the region (Cowx, 1997). Although the establishment of this species has allowed commercial fishing seasons and angler harvest in many countries (Dahl, 1984), it has also disrupted the native fish assemblage in many locations. For example, its introduction into Lake Egridir in Turkey caused the extinction of three species of *Phoxinellus*, of which two were endemic (Crivelli, 1995).

To balance both fishery and conservation objectives, there is a need to develop management plans for *S. lucioperca*. Development of such plans relies on an understanding of size, age structure, and growth characteristics of populations under consideration (Laurec & Le Guen, 1981; Meunier, 1988). The majority of studies on *S. lucioperca* to date have focused on populations located around the Baltic Sea and in Scandinavia (Nagiec, 1977; Lehtonen, 1983; Kangur & Kangur 1996; Hahlbeck & Müller, 2003; Ložys, 2004 and Milardi *et al.*, 2011). In the more southerly regions and/or recently colonized areas, the demographic characteristics of *S. lucioperca* are less well studied (but see Toujani, 1998; M'Hetli, 2001; Poulet, 2004; Argillier *et al.*, 2012; Pérez-Bote and Roso, 2012).

Our objective was to measure the growth of this species in the Ghrib reservoir, Algeria, a location outside the native range of the species. Our results can be used to inform stock assessment analyses for this area.

MATERIALS AND METHODS

STUDY SITE

The study was conducted in the reservoir upstream of the Ghrib dam, which is located in the south of the wilaya of Ain Defla, 150 km from Algiers. The reservoir has a catchment of 22 456 km². Established on Oued Cheliff (36°13'76.77" N and 2°56'70.01" E), it is delimited at the downstream end by the Boughzoul dam, which acts on it as a settling pond (Fig. 1). Water temperatures in this area fluctuate from 6.5°C in winter to a maximum of 30°C in summer.

FISH SAMPLING

Currently, the fish assemblage in Ghrib reservoir includes eight fresh-water species: *Barbus callensis* (Barbell), *Rutilus rutilus* (Roach), *Carassius carassius* (Crucian carp), *Abramis brama* (Bream), *Cyprinus carpio* (Common carp), *Hypophthalmichthys molitrix* (Silver carp), *Hypophthalmichthys nobilis* (Bighead carp), and *Sander lucioperca* (Pike perch). In December 1990, 8 barbels were captured from the Ghrib reservoir but only one pike perch. In June 1991, 5 barbels were captured (Arab work, unpublished). In 2013, we noted a predominance of carps (10 565 were captured: 1945 in June and 2210 in December), pike perches (2603: 173 in June and 990 in December), and native species such as the barbell (1510: 162 in June and 435 in December). In 2014, we captured 12 060 carps (1870 individuals in June and 1280 in December), 4232 pike perches (564 individuals in June and 635 in December), and 1221 barbells (103 individuals in June and 130 in December).

Samples of *S. lucioperca* were collected monthly from May 2013 to April 2014 in the Ghrib Reservoir by setting gill nets (18, 25, 35, 45 and 65 mm mesh size). The nets were set at the surface in the evening at ~5 p.m., and retrieved the next morning at around 6 a.m., then again at 12 a.m. Fish were immediately transported to the laboratory and measured for total length (TL; to the nearest lower 0.1 cm); fork length (FL; to the nearest lower 0.1 cm); standard length (SL; to the nearest lower 0.1 cm); total weight (TW; to the nearest 0.001 g); eviscerated weight (EW; to the nearest lower 0.001 g), and sex. Several scales (10-20) were removed from the area below the pectoral fin, cleaned with KOH 5% and rinsed with distilled water, then stored dry in individually labelled plastic vials. The scales were held between two blades and observed with a binocular magnifying glass. Readings were made by three independent observers; the scales were used in subsequent analyses only when all three readings were in agreement.

The morphometric relationship between TL or FL and scale-size were of the type $Y = aX + b$. The relationship between length and weight was described by: $W = a \times TL^b$ (Ricker, 1973), where W is the total weight (TW) or the eviscerated weight (EW) in grams, a is the intercept, b is the slope (fish growth rate), and TL is the total length in centimetres. Differences in the relationship between total length and total weight between sexes were tested using ANCOVA. We tested for isometric growth using a t -test. A marginal increment analysis was used to validate annual growth increment formation (Beamish & McFarlane, 1983). The monthly mean of the marginal increment (MI) was calculated using: $MI = (R - R_n) / (R_n - R_{n-1})$. R , R_n , and R_{n-1} are, respectively, the radius, the radius of the last and the next-to-last growth rings. Monthly values of MI were compared using a one-way ANOVA test, followed by a multiple sample comparison of means (Dagnelie, 1975).




Figure 1.— Location of the study site, Ghrif reservoir, Algeria.

The age-length relationship was back-calculated using the Lee method (1920). The relationship between fish length (TL) and total scale radius (R) was obtained by a regression $TL = f(R)$ based on 230 scales ($21 \leq TL \leq 88.2$ cm; $0.57 \leq R \leq 6.24$ mm). Sizes-at-age (age-length key) were compared with the results of the back-calculation. The growth values obtained by scale measurement were analysed and compared with those of the theoretical growth calculated according to the model of Von Bertalanffy (1938) using FiSAT II (version 1.2.2) (Gayanilo *et al.*, 1996). The model was expressed by the equation: $TL = L_{\infty} (1 - e^{-k(t-t_0)})$, where TL is the size of fish at the moment t , L_{∞} is the theoretical maximum size that the fish is likely to reach, and t_0 is theoretical time when $TL = 0$. Sample manpower was classified per month and class of size (5 cm interval). Absolute growth was calculated based on analysis of size distribution modes. The various modes correspond to pseudocohorts of different ages. For this analysis, we used FiSAT II (Gayanilo *et al.*, 1996), which allows estimation of L_{∞} growth, K , and t_0 (Bhattacharya, 1967). These parameters were adjusted using the Von Bertalanffy equation and were compared with those obtained by the direct method of estimating growth. A growth performance index ($\phi' = \log k + 2 \log L_{\infty}$) was calculated to compare the results obtained in this study with results published elsewhere (Pauly & Munro, 1984).

RESULTS

A total of 891 scales were used to assess fish age. Age was assigned for 849 (95.28 %) scales: 375 females, 474 males (Tab. I). The range in total length and total weight for female and male fish was 21.8 - 83.8 cm and 86 - 4935 g, and 21.0 - 88.2 cm and 77 - 7245 g, respectively. The length-length and weight-length relationships are summarized in Table I. The equations expressing the length-weight relationship for pike perch were indicative of positive allometric growth for both sexes (paired t -test, females: t -test = 5.26, $P < 0.05$; males: t -test = 2.99, $P < 0.05$). The allometric coefficient of the regression in females was significantly higher than in males (ANCOVA, $F = 17.89$, $df = 1$, $P < 0.05$).

The equation expressing the linear relationship between total length (TL) and the scale radius (R) was: $TL = 105.06 R + 78.86$ mm, ($r = 0.96$; $P \leq 0.001$). The ordinate to the origin of this equation (78.86 mm) corresponds to the theoretical fish size at the time of first scale formation. Comparison of successive monthly mean marginal increment values for scales using one-way ANOVA ($F = 7.63$; $P < 0.05$) revealed a significant difference between only two consecutive months, February and March (Fig. 2). Thus, the rings were considered to be annual increments.

Minimum values of marginal increment were recorded in March, which is the time of annulus formation.

TABLE I

Morphometric values and length-weight relationships for females, males, and unknown sex of Sander lucioperca in the Ghrib reservoir, Algeria

	N	r	Equation	Sy	Sx	Equation	r
Females	375	0.98	TL = 1.17 SL + 2.361	86.634	73.313	TW = 0.0025 TL ^{3.317}	0.98
	375	0.87	FL = 1.07 SL + 1.871	84.006	73.313	EW = 0.0022 TL ^{3.327}	0.98
	375	0.88	TL = 0.96 FL + 5.688	86.634	84.006	-	-
Males	474	0.99	TL = 1.20 SL + 1.334	72.291	59.841	TW = 0.0047 TL ^{3.142}	0.98
	474	0.97	FL = 1.09 SL + 1.205	66.518	59.841	EW = 0.0042 TL ^{3.147}	0.98
	474	0.97	TL = 1.07 FL + 1.015	72.291	66.518	-	-
Females	849	0.99	TL = 1.19 SL + 1.673	86.370	72.066	TW = 0.0033 TL ^{3.237}	0.98
+	849	0.93	FL = 1.08 SL + 1.667	80.693	72.066	EW = 0.003 TL ^{3.271}	0.98
Males	849	0.93	TL = 1.03 FL + 2.601	86.370	80.693	-	-

TL = overall length (cm); FL = length in the fork (cm); SL = standard length (cm); N = number of individuals; r = coefficient of correlation; Sy = standard deviation y; Sx = standard deviation x; TW = total weight; EW = eviscerated weight (g).




Figure 2.— Monthly evolution of marginal increment (MI) for all *Sander lucioperca* scales sampled from May 2013 to April 2014 at Ghrib reservoir, Algeria. Vertical error bars represent the standard deviation.

The age range of the sampled fish was 1-9 years, with of the majority (75.84 %) of individuals being in the 3-4 year age class (Tab. II). Back-calculated age-length pairs (Tab. III) did not reveal any difference from the observed mean size-at-age determined by direct reading of scales. The median values of the sizes corresponding to each size class were adjusted using the Von Bertalanffy equation: $TL = 125.72 (1 - e^{-0.13 (t + 0.89)})$ with $r = 0.90$ (Fig. 3). Growth performance index (ϕ') was 3.31.

Analysis of the size distribution modes yielded estimates of age and growth (Tab. IV). The median values of the sizes corresponding to each size class were adjusted using a Von Bertalanffy equation: $TL = 132.17 (1 - e^{-0.12 (t + 0.95)})$ with $r = 0.979$ (Fig. 3). There was no significant difference between the results obtained using the two methods (χ^2 calculated = 1.7 < χ^2 theoretical = 14.07; $P = 0.05$).

TABLE II

Age-length key obtained by direct scale reading for *Sander lucioperca* in the Ghrib reservoir, Algeria.

Total length interval (cm)	Age (years)									N
	1	2	3	4	5	6	7	8	9	
20.5 – 22.5	3									3
22.5 – 24.5	19									19
24.5 – 26.5	24									24
26.5 – 28.5	25									25
28.5 – 30.5	3	5								8
30.5 – 32.5		11								11
32.5 – 34.5		18								18
34.5 – 36.5		25	0							25
36.5 – 38.5			45							45
38.5 – 40.5			96							96
40.5 – 42.5			128							128
42.5 – 44.5			81	33						114
44.5 – 46.5				109						109
46.5 – 48.5				47						47
48.5 – 50.5				36						36
50.5 – 52.5				28						28
52.5 – 54.5				29						29
54.5 – 56.5				12	17					29
56.5 – 58.5					22					22
58.5 – 60.5					11					11
60.5 – 62.5					5					5
62.5 – 64.5					0	2				2
64.5 – 66.5						5				5
66.5 – 68.5						0	1			1
68.5 – 70.5							2			2
70.5 – 72.5							0			0
72.5 – 74.5							0			0
74.5 – 76.5							2	0		2
76.5 – 78.5								2		2
78.5 – 80.5								0		0
80.5 – 82.5								0		0
82.5 – 84.5								1		1
84.5 – 86.5								0		0
86.5 – 88.5								0	2	2
N	74	59	350	294	55	7	5	3	2	849
%N	8.71	6.95	41.22	34.62	6.47	0.82	0.58	0.35	0.23	100
TL	M	23.35	30.86	39.10	49.78	59.60	67.83	75.70	83.40	88.10
	E	1.94	1.71	1.68	3.37	1.86	0.94	3.71	4.02	0.14

TL = total length; N = number of fish measured; E = standard deviation; M = mean.

DISCUSSION

We found no difference in relative growth of *S. lucioperca* based on TL/SL, TL/FL, and FL/SL, consistent with Copp *et al.* (2003); in the three types of length-length relationship, χ^2 calculated = $0.99 < \chi^2$ theoretical = 10.82; $p \leq 0.001$. Similarly, Goubier (1975) and Krpo-Cetkovic & Stamenkovic (1996) found no evidence for a difference in growth between the sexes.

TABLE III

Total length (cm) at the deposition of each growth ring in the scales of *Sander lucioperca* from the Ghrif reservoir, Algeria

Age		TL ₁	TL ₂	TL ₃	TL ₄	TL ₅	TL ₆	TL ₇	TL ₈	TL ₉
I	N	74								
	M	21.50								
	E	1.91								
II	N	59	59							
	M	21.30	28.60							
	E	2.14	1.97							
III	N	350	350	350						
	M	21.80	31.30	37.70						
	E	2.10	2.12	1.83						
IV	N	294	294	294	294					
	M	21.90	28.20	39.10	47.60					
	E	2.10	1.95	1.94	3.74					
V	N	55	55	55	55	55				
	M	21.30	28.70	37.60	49.90	57.80				
	E	2.06	1.99	1.88	3.86	2.06				
VI	N	7	7	7	7	7	7			
	M	21.70	28.50	37.80	48.70	59.90	66.80			
	E	2.09	1.97	1.88	3.85	2.14	1.65			
VII	N	5	5	5	5	5	5	5		
	M	21.50	28.60	37.30	48.60	58.80	67.20	74.90		
	E	2.07	1.98	1.86	3.76	2.1	1.66	0.51		
VIII	N	3	3	3	3	3	3	3	3	
	M	21.40	28.80	37.30	48.40	58.70	66.90	74.80	82.90	
	E	2.06	1.99	1.85	3.82	2.09	1.65	0.50	5.01	
VIII	N	2	2	2	2	2	2	2	2	2
	M	21.40	28.60	37.60	48.60	58.80	66.90	74.70	81.70	86.90
	E	2.06	1.98	1.85	3.77	2.0	1.65	0.48	4.13	0.15
Total	N	849	775	716	366	72	17	10	5	2
	M	21.53	28.91	37.77	48.63	58.80	66.95	74.80	82.30	87.70
	E	1.95	2.02	1.86	3.22	1.80	1.40	3.77	2.48	0.17

N = number of fish sampled; M = mean of the length; E = standard deviation. TL₁ to TL₉ are the back-calculated sizes-at-age.




Figure. 3 — Theoretical curves of the growth in length of *Sander lucioperca* from Ghrif reservoir, Algeria. TL = length at the time of growth ring deposition (calculated using Von Bertalanffy model).

Scale marginal increment values suggest that only one growth ring is formed annually, in February. The high percentage of scales with hyaline edges observed at the end of the winter and the beginning of spring may reflect physiological stress during the spawning season (February-May) (Poulet, 2004). These months also constitute the period during which the temperatures in the Ghrif reservoir are the lowest, close to 6.5°C.

TABLE IV

Growth parameters and age of Sander lucioperca from the Ghrib dam, Algeria, estimated by direct (scale method) and indirect (pseudocohort analysis) methods

Parameters	$L_{\infty} = 125.72$ (cm) $k = 0.13$ year ⁻¹ $t_0 = -0.89$ year		$L_{\infty} = 132.17$ (cm) $k = 0.12$ year ⁻¹ $t_0 = -0.95$ year	
Age group	Direct method		Indirect method	
	Sizes observed (cm) (Scalimetry)	Sizes calculated (cm) (Von Bertalanffy)	Sizes observed (cm) (Bhattacharya)	Sizes calculated (cm) (Von Bertalanffy)
I	24.90	23.80	22.57	22.87
II	34.20	32.50	31.90	32.50
III	44.40	43.20	43.20	43.70
IV	56.50	55.40	55.40	56.24
V	66.00	65.10	65.23	65.35
VI	74.20	73.30	73.48	73.67
VII	80.40	79.80	80.17	80.64
VIII	85.80	84.70	86.73	86.93
VIII	87.60	86.40	87.90	88.20

TABLE V

Growth parameters (L_{∞} , k , t_0), growth performance indexes (ϕ'), and parameters for the weight-length relationship (a , b) of Sander lucioperca at different localities

Locality and author	a	b	L_{∞}	k	t_0	ϕ'
Sidi-Salem dam, Tunisia (Toujani, 1998)	-	-	54.44	0.094	-3.346	2.44
Seyhan Dam Lake, Turkey (Özyurt & Avşar, 2002)	-	-	54.44	0.094	-3.346	2.44
German coastal waters, Baltic Sea (Hahlbeck & Müller, 2003)	0.0032	3.283	141.3	0.085	-1.354	3.23
Hirfanli Dam Lake, Turkey (Ablak & Yilmaz, 2004)	10×10^{-6}	3.07	-	-	-	-
Lake İğirdir, Turkey (Balik <i>et al.</i> , 2004)	0.006	3.148	95.4	-0.084	1.563	2.88
Lake Eğirdir, Turkey (İzci & Kuşat, 2006)	0.022	2.742	156.95	-0.045	2.622	3.04
Iranian coastal waters, Caspian Sea (Abdolmalaki & Iwona Psuty, 2007)	-0.0206	2.85	55.05	0.15	-2.59	2.65
Southern Finland (Milardi <i>et al.</i> , 2011)	0.00217	3.371	69.56	0.10	-0.12	2.63
Tunisian reservoirs (M'Hetli <i>et al.</i> , 2011)	5×10^{-6}	3.06	-	-	-	-
French reservoir (Argillier <i>et al.</i> , 2012)	1.91×10^{-6}	5	8	3	38	4
Nature Reserve "Koviljsko-Petrovaradinski Rit", Serbia (Lujčić <i>et al.</i> , 2013)	-	-	49.55	0.25	-0.01	2.78
Lake Marmara, Turkey (Ilhan & San, 2015)	0.0091	2.996	-	-	-	-
Ghrib dam (Present study)	0.0033	3.237	125.72	0.13	-0.89	3.31

Nine age-classes were documented in the current study. This is almost similar to results obtained in the Camargue, France (7 year classes; Poulet, 2004), higher than the numbers estimated for the Sidi-Salem dam, Tunisia (6, Toujani & Kraiem, 2002) and in Turkish reservoirs (5, Ablak & Yilmaz, 2004; Balik *et al.*, 2004), but lower than the numbers in two French reservoirs (14, Argillier *et al.*, 2012) and in Lake Peipsi, Estonia (10, Kangur & Kanguur, 1996) where the species is native compared with the other areas where it is introduced. However, these differences may be the result of insufficient and/or biased sampling methods or to overfishing resulting in smaller size-at-age (Bouhamou *et al.*, 2015).

The exponent of the length-weight relationship (b value) for Ghrib reservoir indicates positive allometric growth ($b = 3.23$) (Tab. V). This value is comparable with those found for populations in Europe, including France ($b = 3.25$) (Argillier *et al.*, 2012) and German coastal waters of the Baltic Sea ($b = 3.28$) (Hahlbeck & Müller, 2003), and in Asia (Turkey: $b = 3.14$; Balik *et al.*, 2004). The allometry coefficients of the length-weight relationship in Turkey ($b = 3.07$, Ablak & Yilmaz, 2004), $b = 2.996$, Ilhan & San, 2015) and in Tunisia ($b = 3.06$, M'Hetli *et al.*, 2011) indicate isometric growth. Interestingly, in Turkey (İzci and Kuşat, 2006) and Iranian coastal waters of the Caspian Sea (Abdolmalaki & IwonaPsuty, 2007) *S. lucioperca* exhibits a completely different pattern of growth (coefficient of allometry $b = 2.74$ and $b = 2.85$, respectively). This

geographical variation in the coefficient of allometry may reflect the degree of acclimatization of *Sander lucioperca* to its location. However, the rate of allometry is known to vary among species, location in time and space (e.g. in relation to hydro-biological conditions, availability of food, sex, and maturity), and the demographic structure of the population (density and thus trophic competition); disease and parasite loads can also affect the value of b (Le Cren 1951; Bagenal & Tesch, 1978).

In Ghrib reservoir, the first annual deceleration of *S. lucioperca* growth occurs at a larger size ($L_1 = 22$ cm) than in the lakes and reservoirs of Europe: Lake Orsjön ($L_1 = 8.3$ cm, Svårdson & Molin, 1973), Lake of Créteil ($L_1 = 9.1$ cm, Gerdeaux, 1986), the Danube, Hungary ($L_1 = 17.8$ cm, Schmid, 1995), Lake Peipsi ($L_1 = 12.3$ cm, Kangur & Kanguur, 1996) and in the “Koviljsko-Petrovaradinski Rit” nature reserve ($L_1 = 15.72$ cm, Lujčić *et al.*, 2013).

The results of the size frequency analysis corroborate those of the scale analysis. The Von Bertalanffy growth parameters obtained in this study and those from other studies are given in Table V. The asymptotic length (L_∞) calculated in this study is relatively high compared to those from other areas: Sidi-Salem dam (Toujani, 1998), Seyhan dam (Özyurt & Avşar 2002), Lake İğirdir (Balik *et al.*, 2004), Caspian Sea (Abdolmalaki & Iwona Psuty, 2007), Southern Finland (Milardi *et al.*, 2011), a French reservoir (Argillier *et al.*, 2012), and the “Koviljsko-Petrovaradinski Rit” nature reserve (Lujčić *et al.*, 2013). However, they are not different from values from the German coastal waters of Baltic Sea (Hahlbeck & Müller, 2003) and Lake Eğirdir in Turkey (İzci & Kuşat, 2006). The growth performance index ($\phi' = 3.31$) was higher in fish from Ghrib reservoir compared to those from other areas, except on the Tunisian Sidi-Salem dam (Toujani, 1998) (Tab. V). We speculate that the range in sizes (L_∞) estimated for the same species reflects the specificity and the ecological characteristics of the environment where the fish lives. For example, water temperature can directly affect fish growth by influencing the physiology of fish (Weatherley & Gill, 1987).

The growth of age 1 and older individuals in Ghrib reservoir appears to be among the most rapid observed for the species (Fig. 3). This exceptional growth may be because it is southernmost population within the range. The optimum temperature for the growth of the *S. lucioperca* is ~28-30°C (Hokanson, 1977; Hilge, 1990) but these temperatures are seldom reached in the high latitudes and, if that happens, it is only over short periods. One thus expects to observe very different growth rates according to the latitude in which the population is located. The positive influence of temperature on the growth of the *S. lucioperca* in the current study is consistent with observations of Ložys (2004) and Heikinheimo *et al.* (2014).

The trophic conditions (quality and quantity of food) are also likely to influence growth. For example, individual zooplanktonivores grow less rapidly than their congeneric piscivores (Mooij *et al.*, 1994; Frankiewicz *et al.*, 1996). In Ghrib reservoir, pike perch preys primarily on roach, which is abundant. Interestingly, no native species (barbel) were found in the stomach of pike perch.

The maximum length of *S. lucioperca* observed in Ghrib reservoir was among the highest recorded, with individuals of > 85.0 cm (Tab. IV). Given the geographical location of the Ghrib population, this is somewhat surprising as Bergmann (Mayr, 1956) suggests that individuals at high latitudes are larger than congeneric individuals at lower latitudes. However this rule is often subject to exceptions (Jonsson, 1991). Other factors such as trophic resources can influence the maximum size. There is still a need for additional work on the growth of the *S. lucioperca* throughout its range to understand the factors affecting its growth.

CONCLUSION

We determined the age and growth parameters of *S. lucioperca* by scalimetry and by analysis of size frequencies. Statistically, there was no significant difference between the results obtained

using the two methods. Thus, either method could be used to obtain growth data for *S. lucioperca*. The exceptional growth observed in *S. lucioperca* in the Ghib reservoir reflects the adaptive capacity of the species and explains, in part, its ability to colonize new environments. The Mediterranean climate, combined with the high thermal optimum for *S. lucioperca* growth, likely largely explains this phenomenon. Our results suggest that the introduction of *S. lucioperca* to the northwest Algerian reservoir has been successful. The tolerance of this introduced species appears to allow self-sustaining populations to persist; however, no study has yet been conducted in Algeria on its possible ecological impact. Nevertheless our results can be used as a starting point for evaluating the population dynamics of *S. lucioperca* in Algeria and estimating its population size.

ACKNOWLEDGEMENTS

We would like to express our thanks and deep gratitude to Jean-Claude Michá, Professor Emeritus at the University of Namur, for his helpful advice on methodology, interpretation, and significance of the analyses, and to two anonymous referees for their comments and suggestions on a previous version of this paper.

REFERENCES

- ABDOLMALAKI, S. & IWONA PSUTY, I. (2007).— The effects of stock enhancement of pikeperch (*Sander lucioperca*) in Iranian coastal waters of the Caspian Sea. *ICES J. Mar. Sci.*, 64: 973-980. DOI:10.1093/icesjms/fsm062.
- ABLAKE, Ö. & YILMAZ, M. (2004).— Growth properties of Pikeperch (*Sander lucioperca* (L., 1758)) living in Hirfanlı Dam Lake. *Turk. J. Vet. Anim. Sci.*, 28: 455-463.
- ARGILLIER, C., BARRAL, M. & IRZ, P. (2012).— Growth and diet of the pikeperch *Sander lucioperca* (L.) in two French reservoirs. *Arch. Pol. Fish.*, 20: 191-200. DOI: 10.2478/v10086-012-0024-0.
- BAGENAL, T.B. & TESCH, F.W. (1978).— Age and growth. Pp 101-136 In: T.B. Bagenal (ed.). *Methods for assessment of fish production in fresh waters*. Blackwell Scientific Publications, Oxford.
- BALIK, Ü., ÇUBUK, H., ÖZKOK, R. & UYSAL, R. (2004).— Size composition, growth characteristics and stock analysis of the Pikeperch, *Sander lucioperca* (L. 1758), population in Lake İğirdir. *Turk. J. Vet. Anim. Sci.*, 28: 715-722.
- BEAMISH, R.J. & MCFARLANE, G.A. (1983).— The forgotten requirements for age validation in fisheries biology. *Trans. Am. Fish. Soc.*, 112: 735-743.
- BHATTACHARYA, C.G. (1967).— A simple method of resolution of a distribution into gaussian components. *Biometrics*, 23: 115-135.
- BOUGHAMOU, N., DERBAL, F. & KARA, M.H. (2015).— Age, growth and reproduction of the black sea bream *Spondyliosoma cantharus* (Linnaeus, 1758) (Sparidae) in the Gulf of Annaba (Algeria). *J. App. Ichthyol.*, 31: 773-779. DOI: 10.1111/jai.12714.
- BÖHLING, P., HUDD, R., LEHTONEN, H., KARAS, P., NEUMAN, E. & THORESSON G. (1991).— Variation in year-class strength of different perch (*Perca fluviatilis*) population in the Baltic sea with special reference to temperature and pollution. *Can. J. Fish. Aquat. Sci.*, 48: 1181-1187. DOI: 10.1139/f91-142.
- CAMPBELL, R.N.B. (1992).— Food of an introduced population of pikeperch, *Stizostedion lucioperca* L., in lake Egirdir, Turkey. *Aquacult. Fish. Manage.*, 23: 71-85.
- COPP, G.H., WESLEY, K.J., KOVAC, V., IVES, M.J. & CARTER, M.G. (2003).— Introduction and establishment of the pikeperch *Stizostedion lucioperca* (L.) in Stanborough Lake (Hertfordshire) and its dispersal in the Thames catchment. *The Lond. Nat.*, 82: 139-153.
- COURNIOU, M., DEHOLLAIN, C. & TEILLAC, P. (2003).— *L'implication des fédérations de pêche dans la gestion du sandre*. Montpellier.
- COWX, I.G. (1997).— L'introduction d'espèces de poissons dans les eaux douces européennes : succès économiques ou désastres écologiques? *Bull. Fr. Pêche Piscic.*, 344/345: 57-77.
- CRIVELLI, A.J. (1995).— Are fish introductions a threat to endemic fresh-water fishes in the Northern Mediterranean Region. *Biol. Cons.*, 72: 311-319.
- DAGNELIE, P. (1975).— *Théorie et méthodes statistiques, Applications agronomiques. Volume. 2. Les méthodes de l'inférence statistique*. 2e ed. Presses agronomiques de Gembloux, Gembloux, Belgique.
- DAHL, J. (1984).— A century of Pikeperch in Denmark. *EIFAC Techn. Pap., Suppl.* 2: 344-352.
- DEELDER, C.L. & WILLEMSSEN, J. (1964).— *Synopsis of biological data on pike-perch Lucioperca lucioperca (Linnaeus) 1758*. FAO Fisheries Synopsis 28, Rome.

- FEHRI-BEDOUI, R., GHARBI, H., KRAIEM, M.M. & EL ABED, A. (2001).— *Particularité écologique et niveau de production d'une lagune tunisienne le lac Ichkeul*. Rapport Commission internationale pour l'exploration scientifique de la Mer Méditerranée 36 : 1.
- FRANKIEWICZ, P., DABROWSKI, K. & ZALEWSKI, M. (1996).— Mechanism of establishing bimodality in a size distribution of age 0 pikeperch, *Stizostedion lucioperca* (L.) in the Sulejow Reservoir, Central Poland. *Ann. Zool. Fenn.*, 33: 321-327.
- GAYANILO, F.C., SPARRE, P. & PAULY, D. (1996).— *The FAO-ICLARM stock assessment tools (FISAT). User's Guide*. FAO Computerized Information Series (Fisheries). No. 6. FAO, Rome.
- GERDEAUX, D. (1986). — *Écologie du gardon (Rutilus rutilus L.) et du sandre (Lucioperca lucioperca L.) dans le lac de Crèteil de 1977 à 1982. Étude de la ligulose du gardon*. Thèse en Sciences Naturelles : Université Pierre et Marie Curie, Paris VI.
- GOUBIER, J. (1975).— *Biogéographie, biométrie et biologie du sandre, Lucioperca lucioperca (L.), Ostéichtyen, Percidé*. Thèse de Doctorat d'État : Université Claude Bernard, Lyon I.
- GOUBIER, J. (1977). — Intérêt piscicole des introductions du sandre *Lucioperca lucioperca* (L.). *B. Fr. Pêche Piscic.*, 265:153-155.
- HAHLBECK, E. & MÜLLER, H. (2003).— Results of the investigations of pike-perch carried out in the German coastal waters of the Baltic Sea from 1992 to 2002. *Inf. Fischwirtsch. Fischereiforsch.*, 50: 71-80.
- HEIKINHEIMO, O., PEKCAN-HEKIMB, Z. & RAITANIEMI, J. (2014).— Spawning stock–recruitment relationship in pikeperch *Sander lucioperca* (L.) in the Baltic Sea, with temperature as an environmental effect. *Fish. Res.*, 155: 1-9.
- HILGE, V. (1990).— Beobachtungen zur Aufzucht von Zandern (*Stizostedion lucioperca* L.) in Labor. *Arch. Fischereiwiss.*, 40: 167-173.
- HOKANSON, K.E.F. (1977).— Temperature requirements of some percids and adaptations to the seasonal temperature cycle. *J. Fish. Res. Board. Can.*, 34: 1524-1550. DOI: 10.1139/f77-217.
- ILHAN, A. & SARI, H.M. (2015).— Length-weight relationships of fish species in Marmara Lake, west Anatolia, Turkey. *Croa. J. Fish.*, 73: 30-32.
- İZCI, L. & KUŞAT, M. (2006).— Some population parameters of Pikeperch (*Sander lucioperca* (L.,1758)) in Lake Eğirdir. *Süleyman Demirel Üniversitesi. Fen Bil. Ens. Derg.*, 10: 167-172.
- JONSSON, B., L'ABÉE-LUND, J.H., HEGGBERGET, T.G., JENSEN, A.J., JOHNSEN, B.O., NAESJE, T.F. & SAETTEM, L.M. (1991).— Longevity, body size, and growth in anadromous brown trout (*Salmo trutta*). *Can. J. Fish. Aquat. Sci.*, 48: 1838-1845. DOI: 10.1139/f91-217.
- KANGUR, A. & KANGUR, P. (1996).— The condition, length and age distribution of pikeperch, *Stizostedion lucioperca* (L.) in Lake Peipsi. *Hydrobiologia*, 338: 179-183. DOI: 10.1007/BF00031722.
- KARA, M.H. (2012).— Freshwater fish diversity in Algeria with emphasis on alien species. *Eur. J. Wildl. Res.*, 58: 243-253. DOI: 10.1007/s10344-011-0570-6.
- KOONCE, J.F., BAGENAL, T.B., CARLINE, R.F., HOKANSON, K.E.F. & NAGIEC, M. (1977).— Factors influencing year-class strength of Percids: a summary and a model of temperature effects. *J. Fish. Res. Board. Can.*, 34: 1900-1909. DOI: 10.1139/f77-256.
- KOTTELAT, M. & FREYHOF, J. (2007).— *Handbook of European freshwater fishes*. Kottelat, Cornol, Switzerland and Freyhof, Berlin, Germany.
- KRAIEM, M.M., BEN HAMZA, C., RAMDANI, M., FATHI, A.A., ABDELZAHAR, H.M.A. & FLOWER, R.J. (2001). — Some observations on the age and growth of thin-lipped grey mullet, *Liza ramada* Risso, 1826 (Pisces, Mugilidae) in three North African wetland lakes: Merja Zerga (Morocco), Garaat Ichkeul (Tunisia) and Edku Lake (Egypt). *Aquat. Ecol.*, 35: 335-345. DOI: 10.1023/A:1011900631096.
- KRPO-CETKOVIC, J. & STAMENKOVIC, S. (1996).— Morphological differentiation of the pikeperch *Stizostedion lucioperca* (L.) population from the Yugoslav part of the Danube. *Ann. Zool. Fenn.*, 33:711-723.
- LAPPALAINEN, J., ERM, V. & LEHTONEN, H. (1995).— Pickperch, *Stizostedion lucioperca* (L.) catch in relation to juvenile density and water temperature in Parnu Bay, Estonia. *Fish. Manage. Ecol.*, 2: 113-120. DOI: 10.1111/j.1365-2400.1995.tb00104.x.
- LAPPALAINEN, J. & LEHTONEN, H. (1997).— Temperature habitats for freshwater fishes in a warming climate. *Boreal Environ. Res.*, 2: 69-84.
- LAUREC, A. & LE GUEN, J.C. (1981).— *Dynamique des populations marines exploitées. Tome I, concepts et modèles*. Centre National pour l'Exploitation des Océans. Rapports scientifiques et techniques 45.
- LE CREN, E.D. (1951).— The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *J. Anim. Ecol.*, 20: 201-219.
- LEE, R.M. (1920).— A review of the methods of age and growth determination in fish by means of scales. *Fishery Invest.*, 2 (4): 2-32.
- LEHTONEN, H. (1983).— Stocks of pike-perch (*Stizostedion lucioperca* L.) and their management in the Archipelago Sea and the Gulf of Finland. *Finn. Fish. Res.*, 5: 1-16.

- LOSSE, G.F., NAU, W. & WINTER, M. (1991).— *Le développement de la pêche en eau douce dans le Nord de la Tunisie*. Coopération technique tuniso-allemande. CGP Ministère de l'agriculture, Tunis.
- LOŽYS, L. (2004).— The growth of pikeperch (*Sander lucioperca* L.) and perch (*Perca fluviatilis* L.) under different water temperature and salinity conditions in the Curonian Lagoon and Lithuanian coastal waters of the Baltic Sea. *Hydrobiologia*, 514: 105-113. DOI: 10.1023/B:hydr.0000018211.26378.b9.
- LUJIĆ, J., KOSTIĆ, D., BJELIĆ-ČABRILO, O., POPOVIĆ, E., MILJANOVIĆ, B., MARINOVIĆ, Z. & MARKOVIĆ, G. (2013).— Ichthyofauna composition and population parameters of fish species from the special nature reserve "Koviljsko-Petrovaradinski Rit" (Vojvodina, Serbia). *Turk. J. Fish. Aquat. Sci.*, 13: 665-673. DOI: 10.4194/1303-2712-v13_4_12.
- MAYR, E. (1956).— Geographical character gradients and climatic adaptation. *Evolution*, 10: 105-108. DOI: 10.2307/2406103.
- MEDDOUR, A., ROUABAH, A., MEDDOUR-BOUDERDA, K., LOUCIF, N., REMILI, A. & KHATAL, Y. (2005).— Expérimentation sur la reproduction artificielle de *Sander lucioperca*, *Hypophthalmichthys molitrix* et *Aristichthys nobilis* en Algérie. *Rev. Sci. Technol.*, C 23: 63-71.
- MEUNIER, F.J. (1988).— Détermination de l'âge individuel chez les Ostéichtyens à l'aide de la squelettechronologie : historique et méthodologie. *Acta. Oecol. Gener.*, 9: 299-329.
- M'HELTI, M. (2001).— *Le sandre Stizostedion lucioperca (L., 1758 Téléostéen, Percidae) poisson allochtone : étude biologique et essai d'optimisation des critères d'élevage*. Thèse de doctorat : Université de Tunis, El Manar.
- M'HELTI, M., BEN KHEMIS, I., HAMZA, N., TURKI, B. & TURKI, O. (2011).— Allometric growth and reproductive biology traits of pikeperch *Sander lucioperca* at the southern edge of its range. *J. Fish Biol.*, 78:567-579. DOI: 10.1111/j.1095-8649.2010.02878.x.
- MILARDI, M., LAPPALAINEN, J., MALINEN, T., VINNI, M. & RUHJÄRVI, J. (2011).— Problems in managing a slow-growing pikeperch (*Sander lucioperca* (L.)) population in Southern Finland. *Knowl. Manag. Aquat. Ecosyst.*, 400, 08. DOI: 10.1051/kmae/2011010.
- MOOIJ, W.M., LAMMENS, E.H.R.R. & VAN DENSEN, W.L.T. (1994).— Growth rate of 0⁺ fish in relation to temperature, body size, and food in shallow eutrophic Lake Tjeukemeer. *Can. J. Fish. Aquat. Sci.*, 51: 516-526. DOI: 10.1139/f94-054.
- NAGIEC, M. (1977).— Pikeperch (*Stizostedion lucioperca*) in its natural habitats in Poland. *J. Fish. Res. Board. Can.*, 34: 1581-1585. DOI: 10.1139/f77-222.
- ÖZYURT, C.E. & AVŞAR, D. (2002).— Seyhan Baraj Gölü'ndeki (Adana) Sudakların (*Sander lucioperca* Bogustkaya & Naseka, 1996) Bazı Biyolojik Özelliklerinin Belirlenmesi. E.Ü. Su Ürünleri Dergisi. E.U. *J. Fish. Aquat. Sci., Cilt/19*: 77-84. <http://jfas.ege.edu.tr>.
- PAULY, D. & MUNRO, J.L. (1984).— Once more on the comparison of growth in fish and invertebrates. *Fishbyte*, 2: 21.
- PÉREZ-BOTE, J.L. & ROSO, R. (2012).— Growth and length-weight relationships of *Sander lucioperca* (Linnaeus, 1758) in the Alcántara Reservoir, south-western Spain: comparison with other water bodies in Eurasia. *J. Appl. Ichthyol.*, 28: 264-268. DOI: 10.1111/j.1439-0426.2011.01918.x.
- POULET, N. (2004).— *Le sandre (Sander lucioperca (L.)): Biologie, comportement et dynamique des populations en Camargue (Bouches du Rhône, France)*. Thèse de Doctorat : Université de Toulouse III, Paul Sabatier.
- RICKER, W.E. (1973).— Linear regressions in fishery research. *J. Fish. Res. Board Can.*, 30: 409-434.
- SCHMID, J. (1995).— Zum Wachstum des Zander in der Donau. *Fischer. & Teichwirt.*, 2: 54.
- SONESTEN, L. (1991).— The biology of pikeperch - A literature review. *Inf. Sotvattenslab., Drottningholm*, 1: 68-71.
- SVÄRDSON, G. & MOLIN, G. (1973).— The impact of climate on Scandinavian populations of the zander (*Stizostedion lucioperca* L.). *Rep. Inst. Freshw. Res., Drottningholm*, 53: 112-139.
- TOUJANI, R. (1998).— *Le sandre Stizostedion lucioperca (L.1758) de la retenue de Sidi-Salem (Tunisie): Biologie et dynamique de population*. Thèse de doctorat: Université Claude Bernard, Lyon I.
- TOUJANI, R. & KRAIEM, M.M. (2002).— Dynamique de la population de sandre *Stizostedion lucioperca* (L., 1758 pisces, percidae) dans la retenue du barrage de Sidi Salem. *Bull. Inst. Natn. Scien. Tech. Mer de Salammbô.*, Volume. 29.
- TRANDAFIRESCU, I.I., GHITESCU, E. & ILIESCU, M. (1979).— Données préliminaires concernant l'élevage des jeunes de sandre (*Stizostedion lucioperca* Linné, 1758) en milieu saumâtre. *Cercet. Mar. Rech. Mar.*, 12: 261-273.
- VERREYCKEN, H., VAN THUYNE, G. & BELPAIRE, C. (2011).— Length-weight relationships of 40 freshwater fish species from two decades of monitoring in Flanders (Belgium). *J. App. Ichthyol.*, 27:1416-1421. DOI: 10.1111/j.1439-0426.2011.01815.x.
- VON BERTALANFFY, L. (1938).— A quantitative theory of organic growth (inquires on growth laws II). *Hum. Biol.*, 10: 181-213.
- WEATHERLEY, A.H. & GILL, H.S. (1987).— *The biology of fish growth*. Academic Press, Orlando, FL, USA.
- WELCOME, R.L. (1988).— *International introductions of inland aquatic species*. FAO, UN, Rome.